



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

character. Catarrhal affections of the respiratory organs are very common, and cases of grippe prevail to some extent.

Respectfully,

H. S. CAMINERO, M. D.,

Sanitary Inspector, U. S. M. H. S.

The SUPERVISING SURGEON-GENERAL,

U. S. Marine-Hospital Service.

JAPAN.

Sanitary report from Yokohama.

YOKOHAMA, November 9, 1897.

SIR: I have the honor to forward herewith my regular report on infectious diseases in Japan for period October 29 to November 8, inclusive. I beg to invite your attention to the continued prevalence of epidemic dysentery, notwithstanding the low temperature of the present season.

I am, sir, very respectfully, your obedient servant,

STUART ELDRIDGE, M. D.,

Sanitary Inspector, U. S. M. H. S.

The SUPERVISING SURGEON-GENERAL,

U. S. Marine-Hospital Service.

December 17, 1897

1390

[Inclosure.]

Report of infectious diseases in Japan, October 29 to November 8, 1897.

Locality.	Cholera.		Dysentery.		Smallpox.	
	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Kioto Fu.....			88	48		
Osaka Fu.....			73	33	1	
Tokyo Fu.....			59	17		
Aichi Ken.....			69	18		
Akita Ken.....			3	1	1	
Awomori Ken.....			291	49		
Chiba Ken.....			119	26	1	
Fukui Ken.....			23	12		
Fukuoka Ken.....			2	2		
Fukushima Ken.....			28	6	1	
Gifu Ken.....			21	8		
Gumma Ken.....			48	20		
Hogo Ken.....			56	15		
Hiroshima Ken.....			142	96		
Ibaraki Ken.....			37	18		
Ishikawa Ken.....			6	4		
Iwate Ken.....			265	62		
Kagawa Ken.....	(a)					
Kagoshima Ken.....			58	7		1
Kanagawa Ken.....			124	26		
Kochi Ken.....			23	4		
Kumamoto Ken.....	(a)					
Miyagi Ken.....			82	30	1	
Miyazaki Ken.....			10	5		
Miye Ken.....			28	16		
Nagano Ken.....			430	156		
Nagasaki Ken.....	(a)					
Nara Ken.....	(a)					
Niigata Ken.....			697	183	5	1
Oita Ken.....			24	11		
Okayama Ken.....			57	22		
Okinawa Ken.....	(a)					
Saga Ken.....			12	4		
Saitama Ken.....			41	19		
Shidzuoka Ken.....			219	76		
Shiga Ken.....			17	7		
Shimane Ken.....			98	38		
Tochigi Ken.....	(a)					
Tokushima Ken.....			12	7		
Tottori Ken.....			46	20		
Toyama Ken.....	(a)					
Wakayama Ken.....			5	2		
Yamagata Ken.....			7	8		
Yamaguchi Ken.....			140	52		
Yamanashi Ken.....			258	86		
Yehime Ken.....	2		116	29	2	
The Hokkaido.....					37	11
Taiwan (Formosa).....	(a)					
Totals.....	2		3, 834	1, 243	49	13

a No report.